

ANTACID ANTIGAS MAXIMUM STRENGTH- aluminum hydroxide, magnesium hydroxide, simethicone liquid

TARGET Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel USP) 800 mg

Magnesium Hydroxide 800 mg

Simethicone 80 mg

Purpose

Antacid

Antacid

Antacid

Uses

- for the relief of
 - acid indigestion
 - heartburn
 - sour stomach
 - upset stomach associated with these symptoms
 - pressure and bloating commonly referred to as gas
 - overindulgence in food and drink

Warnings

Do not take

more than 60 mL in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

symptoms last more than 2 weeks.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- shake well before using
- mL = milliliter
- do not take more than 60 mL in 24 hours or use the maximum dosage for more than 2 weeks
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and older: take 10-20 mL as needed, between or after meals, at bedtime, or as directed by a doctor
- children under 12 years of age: do not use

Other information

- **each 10 mL contains:** magnesium 335 mg, sodium 10 mg
- store between 20-25°C (68-77°F). Avoid freezing

Inactive ingredients

benzyl alcohol, carboxymethylcellulose sodium, flavor, glycerin, microcrystalline cellulose, purified water, sodium carbonate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-800-910-6874

Principal Display Panel

Compare to active ingredients in Mylanta® Maximum Strength*

Maximum Strength fast acting

antacid

anti-gas

aluminum hydroxide

magnesium hydroxide

simethicone

soothing relief of heartburn,

acid indigestion, sour stomach

sugar, paraben and alcohol free

*This product is not manufactured or distributed by McNeil Consumer Pharmaceuticals Co., distributor of Mylanta® Maximum Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

Distributed by Target Corporation

Minneapolis, MN 55403

Product Label

NDC 11673-528-12

Compare to active ingredients in
Mylanta® Maximum Strength*

maximum strength
fast acting
**antacid
anti-gas**
aluminum hydroxide
magnesium hydroxide
simethicone

soothing relief of heartburn,
acid indigestion, sour stomach
sugar, paraben and alcohol free

up&up

12 FL OZ (355 mL)

PLD-A460B LB005153

Drug Facts

Active ingredients (in each 10 mL) **Purposes**
Aluminum hydroxide (equiv. to dried gel USP) 800 mg.....Antacid
Magnesium hydroxide 800 mg.....Antacid
Simethicone 80 mg.....Antigas

Uses ■ for the relief of
■ acid indigestion
■ heartburn
■ sour stomach
■ upset stomach associated with these symptoms
■ pressure and bloating commonly referred to as gas
■ overindulgence in food and drink

Warnings
Do not take more than 60 mL in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician.
Ask a doctor before use if you have
■ kidney disease
■ a magnesium-restricted diet

*This product is not manufactured or distributed by McNeil Consumer Pharmaceuticals Co., distributor of Mylanta® Maximum Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

Distributed by Target Corporation
Minneapolis, MN 55403
TM & ©2018 Target Brands, Inc.

PLD-A460B LB005154
245 05 0322 R00 C-000703-01-023



PEEL CORNER TO READ COMPLETE DRUG FACTS INFORMATION

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ shake well before using
■ mL = milliliter
■ do not take more than 60 mL in 24 hours or use the maximum dosage for more than 2 weeks
■ measure only with dosing cup provided. Do not use any other dosing device.
■ keep dosing cup with product
■ adults and children 12 years of age and older: take 10-20 mL as needed, between or after meals, at bedtime, or as directed by a doctor
■ children under 12 years of age: do not use

Other information
■ each 10 mL contains: magnesium 335 mg, sodium 10 mg
■ store between 20-25°C (68-77°F). Avoid freezing.

Inactive ingredients benzyl alcohol, carboxymethylcellulose sodium, flavor, glycerin, microcrystalline cellulose, purified water, sodium carbonate, sorbitol, sucralose, xanthan gum

Questions or comments?
Call 1-800-910-6874

TARGET Maximum Strength Antacid anti-gas

ANTACID ANTIGAS MAXIMUM STRENGTH

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-528
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDRO XIDE	800 mg in 10 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDRO XIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
WATER (UNII: 059QF0KO0R)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-528-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2018	12/30/2022

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part331	01/31/2018	12/30/2022

Labeler - TARGET Corporation (006961700)

Revised: 11/2020

TARGET Corporation